

Chapter 13

“Ultra-Processed Foods and Environmental Pollutants: Emerging Determinants of Gut and Mental Health”

Introduction

The global dietary transition towards ultra-processed foods (UPFs), coupled with growing exposure to environmental pollutants such as per- and polyfluoroalkyl substances (PFAS), heavy metals, and microplastics, represents a dual burden on human health. These exposures converge on a common pathway—disruption of gut microbiome composition and intestinal barrier integrity. A compromised gut not only increases systemic inflammation but also directly affects the gut-brain axis, with profound implications for mental health disorders including depression, anxiety, and cognitive decline. Understanding these links is crucial to framing nutrition and environmental safety within a public health perspective.

13.1 Environmental Pollutants and Gut

Integrity Per and Polyfluoroalkyl Substances (PFAS)

PFAS, widely used in food packaging, nonstick cookware, and industrial products, are persistent organic pollutants with bioaccumulative potential.

- **Mechanism:** PFAS alter bile acid metabolism, disrupt epithelial tight junction proteins, and modify gut microbiota, increasing intestinal permeability (7).
- **Health Link:** PFAS exposure correlates with metabolic syndrome, immune dysregulation, and depression (8).

Heavy Metals (Lead, Mercury, Cadmium, Arsenic)

Heavy metals enter food chains via contaminated water, soil, and crops.

- **Mechanism:** Metals generate oxidative stress and mitochondrial dysfunction in intestinal epithelial cells, damage tight junctions, and impair beneficial microbes (9).

- **Health Link:** Chronic exposure is associated with neurotoxicity, mood disturbances, and impaired cognitive function through gut-mediated systemic inflammation (10).

Microplastics

Microplastics are now ubiquitous in food, water, and air.

- **Mechanism:** They act as physical irritants to the gut lining, induce ROS (reactive oxygen species) production, and alter microbial balance. Additives such as bisphenols and phthalates leach into tissues, further impairing gut and endocrine health (11).

- **Health Link:** Animal studies suggest microplastics impair memory and learning via microbiota disruption, with emerging evidence in humans pointing towards anxiety and depressive symptomatology (12).

13.2 Mechanisms of Gut Impact by UPFs and Environmental Pollutants

1. Disruption of Tight Junctions

- o UPFs (emulsifiers, sugars) and pollutants (PFAS, heavy metals) damage tight junction proteins (occludin, claudins, ZO-1), increasing paracellular permeability.

2. Microbial Dysbiosis

- o UPFs promote pathogenic microbial growth, while pollutants selectively kill commensals and reduce diversity. Dysbiosis leads to endotoxemia.

3. Immune Activation and Inflammation

- o Translocation of bacterial lipopolysaccharides (LPS) triggers systemic immune activation, releasing cytokines (IL-6, TNF- α) that influence brain function.

4. Oxidative Stress and Mitochondrial Dysfunction

- o Heavy metals and microplastics induce ROS, damaging epithelial cells and altering microbiota metabolism.

5. Neuroendocrine Disruption

- o Reduced SCFAs and increased inflammatory mediators impair the vagus nerve signaling and hypothalamic-pituitary-adrenal (HPA) axis regulation.

13.3 Implications for Mental Health

The convergence of UPFs and environmental pollutants leads to a “double-hit” model: weakened gut integrity from dietary patterns is exacerbated by pollutant exposure, intensifying inflammation and neurotoxicity. This synergy underlies rising trends in depression, anxiety, and neurodegenerative conditions in urban populations (13,14).

13.4 Synthesis and Perspectives

The combined burden of UPFs and environmental pollutants on gut integrity underscores a pressing need for multidisciplinary intervention. Nutritional reforms promoting minimally processed diets, regulatory policies limiting pollutant exposure, and innovative gut-targeted therapies (e.g., prebiotics, probiotics, postbiotics) represent promising strategies. Future research should focus on human longitudinal cohorts to establish causal links and design culturally tailored interventions in high-risk populations. Addressing gut health is no longer peripheral but central to preventing mental health disorders in an increasingly toxic and processed world.

13.5 Mechanistic Depth: How UPFs & Pollutants Disrupt Gut Health

Modern diets high in **ultra-processed foods (UPFs)** and environmental exposures like **PFAS, BPA, and microplastics** act through multiple overlapping biological mechanisms.

1. Intestinal Permeability (“Leaky Gut”)

- **Emulsifiers** (e.g., polysorbate-80, carboxymethylcellulose): strip away the **mucus layer**, allowing bacteria to directly contact epithelial cells, increasing **tight junction breakdown** (15)
- **Artificial Sweeteners** (e.g., saccharin, sucralose): impair expression of **occludin and claudin**, key proteins maintaining gut barrier integrity (16).
- **PFAS**: disrupt epithelial barrier by altering lipid bilayer composition, increasing permeability to endotoxins (17).
- **BPA**: directly binds estrogen receptors in gut epithelium, altering **tight junction proteins** (ZO-1, occludin) and facilitating leakiness (18).

- **Microplastics:** physically abrade gut lining, causing micro-lesions and permeability increases (19).

2. Immune Activation & Systemic Inflammation

- **Emulsifiers & UPFs:** induce **low-grade chronic inflammation** by promoting bacterial translocation and LPS (endotoxin) entry into blood (20).

- **PFAS & BPA:** act as **endocrine disruptors**, altering cytokine profiles (\uparrow TNF- α , IL-6, IL-1 β) and impairing immune tolerance (21,21).

- **Microplastics:** trigger **innate immune activation** via recognition as foreign particles, stimulating inflammasome pathways (23).

3. Microbiota Diversity & Dysbiosis

- **UPFs:** low fiber, high refined sugars \rightarrow \downarrow **Bifidobacteria&Lactobacillus**, \uparrow **Firmicutes/Bacteroidetes ratio** linked to obesity and inflammation (24).

- **Artificial Sweeteners:** increase **Proteobacteria** (pathobionts) and reduce SCFA producers, impairing gut resilience (25).

- **PFAS & BPA:** reduce **alpha diversity**, decrease beneficial SCFA-producing genera, and increase opportunistic pathogens (26).

- **Microplastics:** alter gut microbial ecology in animal models, leading to reduced richness and overgrowth of inflammatory taxa (27).

4. Neuroinflammation & Gut-Brain Axis

- UPF-driven dysbiosis \rightarrow \downarrow SCFAs (esp. butyrate), impairing **blood-brain barrier integrity** and increasing neuroinflammation.

- PFAS/BPA exposure \rightarrow altered neurotransmitter metabolism and increased oxidative stress in the CNS(28).

- Microplastics shown in mice to translocate beyond gut \rightarrow detected in **blood and brain tissue**, triggering **microglial activation**

Key Point: Despite acting through different molecular routes, UPFs and pollutants converge on **gut permeability, immune**
The Gut-Brain Connection: How Your Microbiome Affects Mental Health | 80

dysregulation, microbial imbalance, and neuroinflammation, collectively fueling risk of mental health disorders.

13.6 Comparative Evidence: Traditional Diets vs UPF-Heavy Diets

Traditional Diets (Mediterranean, Indian home-cooked, East Asian):

- o High in **whole grains, legumes, fermented foods, polyphenol-rich vegetables, and diverse fibers**.
- o Promote **SCFA production** (butyrate, acetate, propionate), which strengthen gut barrier, regulate immune tolerance, and dampen neuroinflammation (29).
- o Fermented foods (e.g., curd, kimchi, idli, dosa) act as **natural probiotics**, increasing **Lactobacillus** and **Bifidobacteria**.
- **Modern UPF-Heavy Diets** (fast foods, packaged snacks, soft drinks):
 - o High in **refined sugars, emulsifiers, additives, low fiber**, and exposed to **plasticizers and PFAS** from packaging.
 - o Promote **gut dysbiosis** (loss of diversity, overgrowth of Proteobacteria), **metabolic endotoxemia**, and **leaky gut syndrome** (30).
 - o Associated with **higher rates of depression, anxiety, and cognitive decline** compared to traditional diets in longitudinal studies (31).

Where traditional diets foster **microbial diversity and gut resilience**, UPF-heavy diets synergize with pollutant exposure to fuel **gut inflammation and mental health vulnerability**.

References

1. Monteiro CA, Cannon G, Moubarac JC, Levy RB, Louzada MLC, Jaime PC. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr.* 2018;21(1):5-17.
2. Zinöcker MK, Lindseth IA. The Western diet-microbiome-host interaction and its role in metabolic disease. *Nutrients.* 2018;10(3):365.
3. Chassaing B, Koren O, Goodrich JK, Poole AC, Srinivasan

S, Ley RE, Gewirtz AT. Dietary emulsifiers impact the mouse gut microbiota promoting colitis and metabolic syndrome. *Nature*. 2015;519(7541):92-6.

4. Suez J, Korem T, Zeevi D, Zilberman-Schapira G, Thaiss CA, Maza O, et al. Artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature*. 2014;514(7521):181-6.

5. DiNicolantonio JJ, Lucan SC, O'Keefe JH. The evidence for fructose harmfulness: A critical review. *BMJ Open Heart*. 2017;4(2):e000617.

6. Silva YP, Bernardi A, Frozza RL. The role of short-chain fatty acids from gut microbiota in gut-brain communication. *Front Endocrinol*. 2020;11:25.

7. Blake BE, Fenton SE. Early life exposure to per- and polyfluoroalkyl substances (PFAS) and latent health outcomes: A review including the placenta as a target tissue and possible driver of developmental origins of health and disease. *Hum Reprod Update*. 2020;26(4):537-57.

8. Sunderland EM, Hu XC, Dassuncao C, Tokranov AK, Wagner CC, Allen JG. A review of the pathways of human exposure to poly- and perfluoroalkyl substances (PFASs) and present understanding of health effects. *J Expo Sci Environ Epidemiol*. 2019;29(2):131-47.

9. Breton J, Massart S, Vandamme P, De Brandt E, Pot B, Foligné B. Ecotoxicology inside the gut: Impact of heavy metals on the mouse microbiome. *BMC Pharmacol Toxicol*. 2013;14:62.

10. Sanders AP, Claus Henn B, Wright RO. Perinatal and childhood exposure to cadmium, manganese, and metal mixtures and effects on cognition and behavior: A review of recent literature. *Curr Environ Health Rep*. 2015;2(3):284-94.

11. Lu L, Luo T, Zhao Y, Cai C, Fu Z, Jin Y. Interaction between microplastics and microbiota: A consideration on environmental animal and human health. *Sci Total Environ*. 2019;667:94-100.

12. Yong CQY, Valiyaveetil S, Tang BL. Toxicity of microplastics and nanoplastics in mammalian systems. *Int J Environ Res Public Health*. 2020;17(5):1509.

13. Lane MM, Davis JA, Beattie S, Gómez-Donoso C, Loughman A, O'Neil A, et al. Ultraprocessed food and chronic noncommunicable diseases: A systematic review and meta-analysis of 43 observational studies. *Obes Rev*. 2021;22(3):e13146.

14. Wu S, Ruan Q, Li B, Wang F, Lin W. The role of environmental pollutants in gut microbiota dysbiosis: A review. *Ecotoxicol Environ Saf.* 2022;239:113626.
15. Chassaing B, et al. Dietary emulsifiers impact the mouse gut microbiota promoting colitis and metabolic syndrome. *Nature.* 2015;519(7541):92–6.
16. Suez J, et al. Artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature.* 2014;514(7521):181–6.
17. Xie G, et al. Exposure to PFAS induces intestinal barrier dysfunction in mice. *Environ Pollut.* 2022;306:119424.
18. Liu Y, et al. Bisphenol A disrupts epithelial barrier integrity via estrogen receptor-mediated pathways. *Toxicol Sci.* 2014;138(1):109–19.
19. Jin Y, et al. Microplastics exposure induced intestinal barrier dysfunction and microbiota dysbiosis in mice. *Sci Total Environ.* 2019;667:445–53.
20. Chassaing B, Gewirtz AT. Gut microbiota, low-grade inflammation, and metabolic syndrome. *Toxicol Pathol.* 2014;42(1):49–53.
21. Grandjean P, et al. Immunotoxic effects of perfluorinated compounds. *Environ Health Perspect.* 2008;116(6):712–8.
22. Rochester JR. Bisphenol A and human health: a review of the literature. *Reprod Toxicol.* 2013;42:132–55.
23. Li B, et al. Microplastics and nanoplastics: new contaminants in food and environment. *TrAC Trends Analyt Chem.* 2020;125:115821.
24. De Filippis F, et al. High-level adherence to a Mediterranean diet beneficially impacts the gut microbiome. *Gut.* 2016;65(11):1812–21.
25. Zhou Z, et al. PFAS exposure alters gut microbiota composition and metabolism. *Sci Total Environ.* 2022;806:150468.
26. Lu L, et al. Uptake and accumulation of polystyrene microplastics in zebrafish and its toxic effects. *Environ Sci Technol.* 2016;50(7):4054–60.
27. Grandjean P, Landrigan PJ. Neurobehavioural effects of developmental toxicity. *Lancet Neurol.* 2014;13(3):330–8.
28. Prüst M, et al. Translocation of microplastics to brain tissue and neurotoxicity: experimental evidence. *Toxicol Lett.* 2020;319:1–8.

29. Nagpal R, et al. Fermented foods, microbiota, and mental health: ancient practice meets modern science. *Front Nutr.* 2018;5:61.
30. Monteiro CA, et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr.* 2019;22(5):936–41.
31. Adjibade M, et al. Ultra-processed food consumption and risk of depressive symptoms: longitudinal study. *Nutrients.* 2019;11(8):1801.