

## Chapter 14

### Sleep, Circadian Rhythms, and the Gut Microbiome – Interconnected Systems in Health and Disease

Dr. Indira Prasad

Additional Professor, Dept of Obstetrics and Gynaecology, AIIMS  
Patna

#### Introduction

Sleep and circadian rhythms coordinate a multitude of physiological processes, from metabolism to immune function. Meanwhile, the gut microbiome – the complex community of microorganisms inhabiting the gastrointestinal tract – has emerged as a major regulator of host health. Growing evidence indicates that these systems do not operate independently: rather, there is a tight, bidirectional interplay among sleep/circadian regulation, gut microbial rhythmicity, and host health outcomes. In this chapter, we synthesize current knowledge on how sleep and circadian rhythms influence gut microbiota oscillations, how disruptions (e.g., jet lag, shift work) may precipitate dysbiosis, and how such dysregulation may affect mental and physical health.

This chapter synthesizes current evidence on the connections between sleep, circadian rhythms, and the microbiome, explains mechanisms linking circadian disruption to gut dysbiosis, and reviews the mental health consequences of microbiome-circadian misalignment. Clinical and public health implications are discussed, as well as emerging interventions.

#### 14.1 The Human Circadian System: An Overview

##### *Central and Peripheral Clocks*

The circadian system is governed by a central pacemaker located in the **suprachiasmatic nucleus (SCN)** of the hypothalamus. The SCN synchronizes peripheral clocks in nearly every tissue – including the gastrointestinal tract, liver, pancreas, and immune cells – via hormonal cues, autonomic signals, feeding rhythms, and temperature cycles.

The molecular clock mechanism is based on transcription-translation feedback loops involving the following gene families:

- **CLOCK** and **BMAL1** (positive regulators)
- **PER1/2** and **CRY1/2** (negative regulators)
- Additional modulators such as **REV-ERB $\alpha$** , **ROR $\alpha$** , and nuclear receptors

These core genes coordinate diurnal oscillations in metabolism, detoxification, immune responses, digestive enzyme secretion, and epithelial cell turnover.

### 14.2 Circadian Control of Gastrointestinal Physiology

The GI system displays strong circadian patterns:

Physiological Feature	Daytime Pattern	Nig
Gastric emptying	Faster	Slower
Intestinal motility	Increased	Decrease
pH variability	More dynamic	Stabilize
Mucus secretion	Higher	Lower
Epithelium proliferation	Peaks during night	Declines

### 14.3 Microbiome Rhythmicity: Diurnal Oscillations in the Gut Ecosystem

#### *Microbial Rhythmic Patterns*

Gut bacteria are not static; many taxa exhibit **predictable oscillations** in abundance, gene expression, and metabolite production. Examples include:

- **Proteobacteria** increased during active phase
- **Firmicutes** enriched during rest/fasting phase
- Microbial functions (carbohydrate metabolism, DNA repair, protein synthesis) oscillate with the feeding cycle

Reitmeier et al. identified arrhythmic microbial signatures associated with increased risk of type 2 diabetes, highlighting the clinical relevance of microbiota rhythmicity (1).

#### 14.4 Drivers of Microbial Oscillations

Microbial rhythms depend on:

- **Feeding–fasting cycles** (primary zeitgeber for the gut/main synchronizing signal)
- **Host metabolic hormones** (insulin, glucagon, GLP-1)
- **Epithelial turnover** timed by circadian genes
- **Bile acid secretion**, which follows circadian patterns
- **Melatonin**, present in the GI tract in higher concentrations than the pineal gland

Circadian clock gene knockout models (e.g., *Bmal1*-null mice) lose microbial rhythmicity entirely (2).

#### 14.5 Feedback: Microbes Influence Host Clock Genes

Microbial metabolites—such as short-chain fatty acids (SCFAs), secondary bile acids, tryptophan metabolites, and neurotransmitter precursors—feed back to modulate:

- Intestinal epithelial clock gene expression
- Hepatic peripheral clocks
- Energy regulation
- Sleep pressure (e.g., via SCFAs influencing vagal pathways)

Bidirectionality is central: **the host sets the rhythm, and microbes fine-tune it.**

#### 14.6 Sleep as a Regulator of the Gut Microbiome

##### *Physiological Effects of Sleep on Gut Microbial Homeostasis*

Sleep ensures coherence among peripheral clocks. During consolidated sleep:

- GI motility slows
- Mucosal immunity synchronizes
- Metabolic pathways shift to fasting state

- Intestinal barrier integrity is strengthened
- Microbial communities enter predictable metabolic phases

Studies show that **even a single night of sleep restriction alters intestinal microbiota composition** and increases pro-inflammatory microbial species (3).

Sleep Fragmentation, Short Sleep, and Dysbiosis

Short sleep (<6 hours) and fragmented sleep are associated with:

- ↓ microbial diversity
- ↑ Firmicutes/Bacteroidetes ratio
- ↑ inflammatory genera (*Ruminococcus*, *Blautia*)
- Altered SCFA profiles
- Increased gut permeability

Mechanistically:

- Sleep loss activates the HPA axis → ↑ cortisol → changes in GI motility and immune regulation
- Sympathetic overactivity alters nutrient absorption and epithelial turnover
- Circadian misalignment disrupts feeding behavior, often leading to late-night eating

These changes destabilize the microbial ecosystem.




Figure 1: Circadian-Microbiome-Brain Axis

## 14.7 Circadian Disruption: Jet Lag, Shift Work, and the Microbiome

### Jet Lag and Social Jet Lag

Jet lag rapidly shifts the central clock, while peripheral clocks (e.g., gut, liver) shift gradually. This leads to **temporary desynchronization**.

Studies show:

- A single 6-hour shift alters microbial functional profiles
- Loss of rhythmicity in carbohydrate and lipid metabolism genes
- Transient increases in **Proteobacteria**, a marker of dysbiosis (4)

Even weekend “social jet lag” – a >2-hour discrepancy between weekday and weekend sleep – has been associated with altered microbial profiles in adolescents and adults.

### *Shift Work: A Model of Chronic Circadian Misalignment*

Shift work exposes individuals to long-term circadian disruption due to:

- Irregular sleep schedules
- Nocturnal eating
- Mismatched light exposure
- Altered activity rhythms

A systematic review reported:

- Decreased  $\alpha$ -diversity
- Higher abundance of *Escherichia/Shigella*, *Blautia*, *Dialister*
- Increased inflammatory metabolic pathways (5)

## 14.8 GI and Metabolic Consequences in Shift Workers

Long-term shift work is associated with:

- IBS-like symptoms
- GERD

- Functional dyspepsia
- Metabolic syndrome
- Increased cardiovascular risk
- Non-alcoholic fatty liver disease

Animal studies mirror these findings: chronic circadian disruption in mice leads to dysbiosis, visceral hypersensitivity, and altered metabolomic profiles consistent with IBS (6).

## 14.9 Mechanisms Linking Circadian Disruption to Dysbiosis

### *Irregular Feeding Patterns and Chrononutrition*

Feeding time is the strongest synchronizer of the gut clock. Circadian misalignment (e.g., eating at night) causes:

- Loss of microbial rhythmicity
- Increased endotoxin production (LPS)
- Metabolic inflexibility
- Increased adiposity

Time-restricted feeding has been shown to restore microbial oscillations even when sleep patterns remain irregular.

### *Hormonal and Immune Disruption*

Circadian disruption alters secretion of:

- **Cortisol** (disrupted diurnal curve)
- **Melatonin** (delayed or blunted release)
- **Insulin** and postprandial hormone rhythms

These changes modify gut motility, mucus secretion, and barrier function.

- Loss of Microbial Metabolite Rhythmicity
- Metabolites such as:
  - SCFAs (butyrate, acetate)
  - Secondary bile acids
  - Indoles
  - Serotonin precursors

display diurnal patterns. Loss of these rhythms contributes to:

- impaired epigenetic regulation
- metabolic syndrome
- neurobehavioral changes

#### **14.10 Mental Health Implications**

Circadian disruption is strongly linked to mood disorders, and recent evidence suggests the gut microbiome plays a mediating role.

##### *Microbiome–Gut–Brain Axis*

Key pathways include:

- Vagal signaling
- HPA axis modulation
- **Tryptophan–serotonin pathway**
- Inflammatory cytokines
- Neuroactive metabolites

Disrupted microbial rhythms can reduce the production of metabolites essential for mood stabilization.

##### *14.11 Depression, Anxiety, and Circadian/Microbiome Disturbance*

Studies show:

- Social jet lag correlates with higher depressive symptoms
- Shift workers have higher rates of anxiety, depression, and cognitive impairment
- Microbial dysbiosis (e.g., low SCFA-producing bacteria) is associated with major depressive disorder

Milota et al. proposed that circadian disruption combined with dysbiosis reduces resilience of stress pathways, increasing psychiatric vulnerability (7).

##### *14.12 Sleep Disorders and Microbiome Alterations*

Conditions such as insomnia, OSA, and circadian rhythm sleep-wake disorders exhibit:

- increased inflammatory microbial taxa
- decreased SCFA-producing genera
- altered tryptophan–kynurenine metabolism

This suggests a bidirectional relationship between sleep disturbances and dysbiosis.

### **14.13 Clinical and Public Health Relevance**

#### ***Implications for Healthcare Providers***

Clinicians should recognize that:

- Sleep hygiene influences GI health
- Night-shift healthcare workers face increased metabolic and GI risk
- Gut-directed therapies may aid circadian disorders
- Chrononutrition (meal timing) is relevant to metabolic management

#### ***Health Risks Associated With Circadian–Microbiome***

##### ***Disruption***

- Obesity
- Type 2 diabetes
- IBS
- GERD
- Depression and anxiety
- Cardiovascular disease

##### ***Vulnerable Populations***

- Shift workers (healthcare, aviation, hospitality, manufacturing)
- Frequent travelers
- Adolescents with irregular sleep
- Individuals with mental health disorders

## 14.14 Interventions: Restoring Circadian–Microbiome Harmony

### Chronotherapy

- **Light exposure therapy** (morning light for phase advancement)
- **Melatonin supplementation** (for jet lag and delayed sleep phase)
- Regular sleep schedules

Chronotherapy aligns central and peripheral clocks.

### Chrononutrition

Key evidence-based principles:

- Consistent meal timing
- Avoiding late-night eating
- Time-restricted feeding (8–12 h window)
- High-fiber diets supporting rhythmic SCFA production

### *Microbiome-Targeted Therapies*

- Probiotics (e.g., *Bifidobacterium*, *Lactobacillus*)
- Prebiotics (inulin, resistant starch)
- Polyphenol-rich diets (berries, tea, cocoa)
- Fecal microbiota transplantation (emerging, experimental)

### *Behavioral and Lifestyle Interventions*

- Sleep hygiene education
- Strategic napping for shift workers
- Minimizing rotating shift schedules
- Structured exercise (also has circadian effects)

### Challenges and Future Directions

#### *Research Gaps*

- Human trials remain limited; many are cross-sectional
- Differences in sequencing methods complicate comparison

- Inter-individual variability in microbial responses
- Need for long-term interventional studies

### ***Emerging Research Areas***

- Personalized chrononutrition based on microbiome profiling
- Microbiome-derived psychobiotics
- Use of metabolomics to track circadian health
- Artificial intelligence for circadian risk prediction

### ***The Future of Circadian–Microbiome Medicine***

Possible future clinical tools:

- Microbial rhythmicity tests
- Circadian-metabolite-based sleep therapeutics
- Workplace policies to reduce chronodisruption

### **Conclusion**

Sleep, circadian rhythms, and the gut microbiome represent a deeply integrated triad essential for maintaining metabolic, gastrointestinal, immunologic, and neurobehavioral health. Disruptions such as jet lag and shift work can destabilize this network, leading to gut dysbiosis, systemic inflammation, altered neuroendocrine balance, and increased susceptibility to metabolic and mental health disorders.

Understanding these interactions offers new avenues for prevention and treatment—including chronotherapy, chrononutrition, and targeted microbiome interventions. As modern society continues to challenge natural circadian rhythms, incorporating circadian–microbiome concepts into clinical practice and public health strategies is increasingly essential.

### **References**

1. Reitmeier S, et al. Arrhythmic gut microbiome signatures predict risk of type 2 diabetes. *Nature*. 2022.
2. Thaïss CA, et al. Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell*. 2016.

3. Benedict C, et al. *Acute sleep deprivation increases food intake and modifies gut microbiota in humans.* *Nutr Diabetes.* 2016.
4. Voigt RM, et al. *The circadian clock and the microbiome—implications for health and disease.* *Gastroenterology.* 2016.
5. Maki Y, et al. *How shift work affects the gut microbiota: Implications for gastrointestinal diseases.* *Medicina (Kaunas).* 2024.
6. Porcu A, et al. *Chronic circadian disruption induces IBS-like symptoms in mice through gut dysbiosis.* *BMC Microbiol.* 2025.
7. Milota A, et al. *Circadian disruption and mental health: Microbiome-mediated pathways.* *Int J Mol Sci.* 2024.