

Chapter 15

Microbiome and Gender Differences

Dr. Ruchita

Consultant – Infertility, Indira IVF, Kanpur

Introduction

Sex differences in the human microbiome are increasingly recognized and reflect the influence of sex hormones on microbial composition and function. These interactions, termed the “microgenderome,” affect metabolism, immunity, and disease risk across life stages. Hormonal transitions – puberty, PCOS, and menopause – significantly shape the microbiome, while the microbiome modulates sex hormone availability, particularly estrogens.

15.1 Hormonal Transitions and the Microbiome

Puberty:

- Gonadal steroid surges during puberty create sexual dimorphism in gut microbiota.
- Females show enrichment of pathways for carbohydrate and lipid metabolism; males exhibit distinct microbial patterns influenced by testosterone.
- Animal studies confirm sex hormones causally shape microbiota. [1,2]

PCOS:

- Hyperandrogenism in PCOS is linked to gut dysbiosis, including reduced *Lactobacilli* and *Bifidobacteria*, altered *Prevotellaceae* abundance, and decreased α -diversity.
- Dysbiosis may contribute to insulin resistance, inflammation, and hormonal imbalances. [3,4]

Menopause:

- Decline in ovarian hormones leads to reduced microbial diversity and altered Firmicutes/Bacteroidetes ratio.
- These changes can impair the estrobolome, affecting estrogen reabsorption and contributing to metabolic and systemic disease risk. [5,6]

Estrogen-Microbiome Interaction

- Gut microbes express enzymes (β -glucuronidase, hydroxysteroid dehydrogenases) that deconjugate estrogens, enabling enterohepatic recycling.
- Microbiome composition influences circulating estrogen levels, while estrogens shape microbial diversity and abundance.
- Dysbiosis may impair estrogen metabolism, potentially affecting bone, metabolic, and cardiovascular health. [7,8]

Clinical Implications

- Sex hormone-microbiome interactions contribute to metabolic, reproductive, and systemic disease risk.
 - Understanding these dynamics can inform interventions in PCOS, menopause, and other hormone-related conditions.
 - Microbiome-targeted therapies (probiotics, diet, prebiotics) hold potential but require further study. [2,4,7]
- Here's a **concise reference table** summarizing key hormones, life stages, and microbiome effects:

Life Stage/Condition	Key Hormones	Microbiome Changes	Notes / Clinical Implications
Pre-puberty	Low sex steroids	Similar microbiome between sexes	Baseline microbial composition before hormonal influence
Puberty	\uparrow Estrogen (females), \uparrow Testosterone (males)	Sexual dimorphism emerges; females: \uparrow carbohydrate/lipid metabolism pathways; males: distinct microbial composition	Hormones drive gut microbiome differences that may influence metabolism

PCOS	↑ Androgens / estrogen altered	↓ Lactobacilli, Bifidobacteria; altered Prevotellaceae; ↓ α-diversity	Dysbiosis may contribute to insulin resistance, inflammation, and hormonal imbalance
Reproductive	Cyclic	Fluctuating	Microbiome

Pre-menopausal women	estrogen & progesterone	microbiome diversity with menstrual cycle; enriched estrobolome function	can metabolize estrogens, modulating systemic hormone levels
Pregnancy	↑ Estrogen, ↑ Progesterone	↑ Proteobacteria & Actinobacteria; ↓ diversity in late pregnancy	Microbiome supports metabolic adaptation and immune tolerance
Perimenopausal use/ Menopause	↓ Estrogen, ↓ Progesterone	↓ Diversity; ↑ Firmicutes/Bacteroidetes ratio; altered estrobolome	May contribute to metabolic syndrome, bone loss, and systemic inflammation
Postmenopausal use	Low sex steroids	Persistently altered microbiome; impaired estrogen metabolism	Increased risk for metabolic, cardiovascular, and bone-related disorders

Notes:

- “Estrobolome” refers to gut microbial genes capable of metabolizing estrogens.
- Microbiome-hormone interactions are bidirectional: hormones shape microbiota, microbiota modulate hormone availability.
- Clinical relevance: dysbiosis at these stages may contribute to disease risk and could be targeted via diet, probiotics, or other interventions.

Conclusion

Hormonal transitions across life stages drive sex-specific microbiome changes, while the microbiome reciprocally regulates sex hormone availability. The bidirectional interplay between hormones and microbes underlies health and disease susceptibility in a sex-dependent manner, highlighting the microgenderome as a promising focus for personalized medicine.

References

- Sakong I, Park HJ, Lee KY. Sexual dimorphism of gut microbiota at different pubertal status. *Microb Cell Fact*. 2020;19(1):34.
- Santos-Marcos JA, Andersen E, Pérez-Jiménez J, et al. Interaction between gut microbiota and sex hormones and their relation to sexual dimorphism in metabolic diseases. *Biol Sex Differ*. 2023;14(1):4.
- Torres PJ, Ho BS, Diela Z, et al. Gut microbiota in polycystic ovary syndrome (PCOS). *J Clin Endocrinol Metab*. 2018;103(7):2552–2561.
- Zeng B, Li G, Yuan J, et al. Dysbiosis of gut microbiota in polycystic ovary syndrome (PCOS): Evidence from clinical and preclinical studies. *J Transl Med*. 2025;23:??.
- Cross TW, et al. Ovarian hormone loss, gut microbiome alterations, and metabolic disease risk. *Gut Microbes*. 2023.
- Chen I, Wu Z, Harlow SD, et al. Variance as a predictor of health outcomes: Sex hormones and body fat changes in peri- and post-menopausal women. *arXiv*. 2022.
- Baker JM, Al-Nakkash L, Herbst-Kralovetz MM. The vagina-microbiome-estrogen connection: Mechanistic insights and clinical implications. *Biochim Biophys Acta Mol Basis Dis*. 2017;1863(10 Pt A):2449–2459.
- Niccolò P, Federica M, Francesca R, et al. Gut microbiota and sexual hormones: The “microgenderome” perspective. 2021.